

MINIMAX MODULES WHIT RESPECT TO AN IDEAL AND EXTENSION FUNCTORS OF LOCAL COHOMOLOGY MODULES

B. VAKILI

ABSTRACT. Let R be a commutative Noetherian ring and \mathfrak{a} an ideal of R . The concepts of \mathfrak{a} -minimax and \mathfrak{a} -cominimax modules were introduced by Azami, Naghipour and Vakili in [1] as generalization of minimax and \mathfrak{a} -cofinite modules, respectively. The finiteness of extension functors of local cohomology modules was viewed by Dibaei and Yassemi in [4]. In this paper, we discuss the \mathfrak{a} -minimaxness of extension functors of local cohomology modules, in several cases. Let M be an \mathfrak{a} -minimax R -module, and let t be a non-negative integer such that $H_{\mathfrak{a}}^i(M)$ is \mathfrak{a} -cominimax for all $i < t$. In [1, Theorem 4.1] we have shown that for any \mathfrak{a} -minimax submodule N of $H_{\mathfrak{a}}^t(M)$ and for any finitely generated R -module L with $\text{Supp } L \subseteq V(\mathfrak{a})$, the R -module $\text{Hom}_R(L, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax. In this paper, it is shown that $\text{Ext}_R^i(L, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax for $i = 0, 1$; in addition, if $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(M))$ is \mathfrak{a} -minimax, then $\text{Ext}_R^i(L, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax for $i = 0, 1, 2$.

1. INTRODUCTION

Throughout this paper, R will always be a commutative Noetherian ring with non-zero identity, and \mathfrak{a} will be an ideal of R . Let M be an R -module. The \mathfrak{a} -torsion submodule of M is defined as $\Gamma_{\mathfrak{a}}(M) = \bigcup_{n \geq 1} (0 :_M \mathfrak{a}^n)$. The i^{th} local cohomology functor $H_{\mathfrak{a}}^i(\cdot)$ is defined as the i^{th} right derived functor $\Gamma_{\mathfrak{a}}(\cdot)$. It is known that for each $i \geq 0$ there is a natural isomorphism of R -modules

$$H_{\mathfrak{a}}^i(M) \cong \varinjlim_{n \geq 1} \text{Ext}_R^i(R/\mathfrak{a}^n, M).$$

We refer the reader to [6] or [3] for the basic properties of local cohomology.

An R -module M is said to have *finite Goldie dimension* (written $G \dim M < \infty$), if the injective hull $E(M)$ of M decomposes as a finite direct sum of indecomposable (injective) submodules. It is clear by the definition of the Goldie dimension that

$$G \dim M = \sum_{\mathfrak{p} \in \text{Ass}_R M} \mu^0(\mathfrak{p}, M),$$

where $\mu^0(\mathfrak{p}, M)$ is the 0-th Bass number of M with respect to prime ideal \mathfrak{p} and $\text{Ass}_R M$ is the set of associated prime ideals of R . The \mathfrak{a} -relative Goldie dimension of M has been

Key words and phrases. Goldie dimension, \mathfrak{a} -minimax modules, \mathfrak{a} -cominimax modules, local cohomology.

2000 *Mathematics Subject Classification.* 13D45, 13E05.

The research was supported in part by a grant from Islamic Azad University, Shabestar Branch.
e-mail: bvakil2004@yahoo.com and bvakil@iaushab.ac.ir.

studied by Divaani-Aazar and Esmkhani in [5] and it is defined as

$$G \dim_{\mathfrak{a}} M = \sum_{\mathfrak{p} \in V(\mathfrak{a})} \mu^0(\mathfrak{p}, M).$$

In particular, they proved that $G \dim_{\mathfrak{a}} M = G \dim \Gamma_{\mathfrak{a}}(M)$. An R -module M is said to have *finite \mathfrak{a} -relative Goldie dimension* if the Goldie dimension of the \mathfrak{a} -torsion submodule $\Gamma_{\mathfrak{a}}(M)$ of M is finite.

In [7], Zöschinger introduced the interesting class of minimax modules. An R -module M is said to be a *minimax* module, if there is a finitely generated submodule N of M , such that M/N is Artinian. Zöschinger, has given in [7] and [8] many equivalent conditions for a module to be minimax. In particular, when R is a Noetherian ring, a module is minimax if and only if each of its quotients has finite Goldie dimension. The concepts of \mathfrak{a} -minimax and \mathfrak{a} -cominimax modules were introduced by Azami, Naghipour and Vakili in [1] as generalization of minimax and \mathfrak{a} -cofinite modules. An R -module M is called *\mathfrak{a} -minimax* if the \mathfrak{a} -relative Goldie dimension of any quotient module of M is finite. It is clear that if M is \mathfrak{a} -torsion then M is \mathfrak{a} -minimax if and only if M is minimax. Also, we say that an R -module M is *\mathfrak{a} -cominimax* if the support of M is contained in $V(\mathfrak{a})$ and $\text{Ext}_R^i(R/\mathfrak{a}, M)$ is \mathfrak{a} -minimax for all $i \geq 0$.

The finiteness of extension functors of local cohomology modules was viewed by Dibaei and Yassemi in [4]. In this paper, we discuss the \mathfrak{a} -minimaxness of extension functors of local cohomology modules, in several cases. Let M be an \mathfrak{a} -minimax R -module, and let t be a non-negative integer such that $H_{\mathfrak{a}}^i(M)$ is \mathfrak{a} -cominimax for all $i < t$. In [1, Theorem 4.1] we have shown that for any \mathfrak{a} -minimax submodule N of $H_{\mathfrak{a}}^t(M)$ and for any finitely generated R -module L with $\text{Supp } L \subseteq V(\mathfrak{a})$, the R -module $\text{Hom}_R(L, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax. This generalizes the main result of Brodmann and Lashgari [2]. In this paper, it is shown that both $\text{Hom}_R(L, H_{\mathfrak{a}}^t(M)/N)$ and $\text{Ext}_R^1(L, H_{\mathfrak{a}}^t(M)/N)$ are \mathfrak{a} -minimax; in addition, if $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(M))$ is \mathfrak{a} -minimax, then $\text{Hom}_R(L, H_{\mathfrak{a}}^t(M)/N)$, $\text{Ext}_R^1(L, H_{\mathfrak{a}}^t(M)/N)$ and $\text{Ext}_R^2(L, H_{\mathfrak{a}}^t(M)/N)$ are all \mathfrak{a} -minimax.

2. THE RESULTS

To prove the main results of this paper, we first bring the following lemma.

Lemma 2.1. *Let M be an R -module and put $L = E(M/\Gamma_{\mathfrak{a}}(M))/M/\Gamma_{\mathfrak{a}}(M)$. Then*

- (i) $H_{\mathfrak{a}}^i(L) \simeq H_{\mathfrak{a}}^{i+1}(M)$ and $\text{Ext}_R^i(R/\mathfrak{a}, L) \simeq \text{Ext}_R^{i+1}(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M))$ for all $i \geq 0$.
- (ii) $\text{Ext}_R^1(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M)) \simeq \text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^1(M))$.

Proof. For (i) See [4, Remark 2.1], and for (ii) See to the proof of [4, Theorem B]. \square

Proposition 2.2. *Let M be an R -module. If $\text{Ext}_R^1(R/\mathfrak{a}, M)$ and $\text{Ext}_R^2(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(M))$ are \mathfrak{a} -minimax, then $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^1(M))$ is \mathfrak{a} -minimax.*

Proof. Consider the exact sequence

$$\text{Ext}_R^1(R/\mathfrak{a}, M) \rightarrow \text{Ext}_R^1(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M)) \rightarrow \text{Ext}_R^2(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(M)).$$

Therefore, it follows from [1, Proposition 2.3] and assumption that $\text{Ext}_R^1(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M))$ is \mathfrak{a} -minimax, and so by previous lemma $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^1(M))$ is \mathfrak{a} -minimax. \square

Now, we are ready to present some sufficient conditions for the \mathfrak{a} -minimaxness of extension functors of local cohomology modules.

Theorem 2.3. *Let M be an R -module. Let t be a non-negative integer such that the local cohomology modules $H_{\mathfrak{a}}^i(M)$ are \mathfrak{a} -cominimax for all $i < t$.*

(i) *If $\text{Ext}_R^{t+1}(R/\mathfrak{a}, M)$ is \mathfrak{a} -minimax, then $\text{Ext}_R^1(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax for any \mathfrak{a} -minimax submodule N of $H_{\mathfrak{a}}^t(M)$.*

(ii) *If $\text{Ext}_R^t(R/\mathfrak{a}, M)$ and $\text{Ext}_R^{t+1}(R/\mathfrak{a}, M)$ are \mathfrak{a} -minimax, then for any \mathfrak{a} -minimax submodule N of $H_{\mathfrak{a}}^t(M)$ and for any finitely generated R -module L with $\text{Supp } L \subseteq V(\mathfrak{a})$, the R -module $\text{Ext}_R^i(L, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax for all $i \leq 1$.*

(iii) *If both $\text{Ext}_R^{t+2}(R/\mathfrak{a}, M)$ and $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(M))$ are \mathfrak{a} -minimax, then the R -module $\text{Ext}_R^2(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax for any \mathfrak{a} -minimax submodule N of $H_{\mathfrak{a}}^t(M)$.*

(iv) *If the R -modules $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(M))$ and $\text{Ext}_R^i(R/\mathfrak{a}, M)$ for $i = t, t+1, t+2$ are \mathfrak{a} -minimax, then for any \mathfrak{a} -minimax submodule N of $H_{\mathfrak{a}}^t(M)$ and for any finitely generated R -module L with $\text{Supp } L \subseteq V(\mathfrak{a})$, the R -module $\text{Ext}_R^i(L, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax for all $i \leq 2$.*

(v) *If $\text{Ext}_R^{t+1}(R/\mathfrak{a}, M)$ and $\text{Ext}_R^2(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M))$ are both \mathfrak{a} -minimax, then the R -module $\text{Hom}_R(L, H_{\mathfrak{a}}^{t+1}(M)/N)$ is \mathfrak{a} -minimax for any \mathfrak{a} -minimax submodule N of $H_{\mathfrak{a}}^{t+1}(M)$ and for any finitely generated R -module L with $\text{Supp } L \subseteq V(\mathfrak{a})$.*

Proof. (i) The exact sequence

$$0 \rightarrow N \rightarrow H_{\mathfrak{a}}^t(M) \rightarrow H_{\mathfrak{a}}^t(M)/N \rightarrow 0$$

induces the following exact sequence,

$$\text{Ext}_R^1(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M)) \rightarrow \text{Ext}_R^1(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M)/N) \rightarrow \text{Ext}_R^2(R/\mathfrak{a}, N).$$

Since $\text{Ext}_R^2(R/\mathfrak{a}, N)$ is \mathfrak{a} -minimax, so in view of [1, Proposition 2.3], it is sufficient to show that $\text{Ext}_R^1(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M))$ is \mathfrak{a} -minimax. We use induction on t . When $t = 0$, then by assumption $\text{Ext}_R^1(R/\mathfrak{a}, M)$ is \mathfrak{a} -minimax, and so the exact sequence

$$0 = \text{Hom}_R(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M)) \rightarrow \text{Ext}_R^1(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(M)) \rightarrow \text{Ext}_R^1(R/\mathfrak{a}, M)$$

implies that $\text{Ext}_R^1(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(M))$ is \mathfrak{a} -minimax. Now suppose, inductively, that $t > 0$ and suppose that the result has been proved for $t - 1$. As $\Gamma_{\mathfrak{a}}(M)$ is \mathfrak{a} -cominimax, it follows that $\text{Ext}_R^i(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(M))$ is \mathfrak{a} -minimax for all $i \geq 0$. Hence by the exact sequence

$$\text{Ext}_R^{t+1}(R/\mathfrak{a}, M) \rightarrow \text{Ext}_R^{t+1}(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M)) \rightarrow \text{Ext}_R^{t+2}(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(M))$$

and assumption, $\text{Ext}_R^{t+1}(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M))$ is \mathfrak{a} -minimax. Also, as $H_{\mathfrak{a}}^0(M/\Gamma_{\mathfrak{a}}(M)) = \Gamma_{\mathfrak{a}}(M/\Gamma_{\mathfrak{a}}(M)) = 0$ and $H_{\mathfrak{a}}^i(M/\Gamma_{\mathfrak{a}}(M)) \simeq H_{\mathfrak{a}}^i(M)$ for all $i > 0$, it follows by assumption that $H_{\mathfrak{a}}^i(M/\Gamma_{\mathfrak{a}}(M))$ is \mathfrak{a} -cominimax for all $i < t$. Therefore, we may assume that M is \mathfrak{a} -torsion free. Let E be an injective envelope of M and put $M_1 = E/M$. Then by lemma 2.1, $\text{Ext}_R^i(R/\mathfrak{a}, M_1) \simeq \text{Ext}_R^{i+1}(R/\mathfrak{a}, M)$ and $H_{\mathfrak{a}}^i(M_1) \simeq H_{\mathfrak{a}}^{i+1}(M)$ for all $i \geq 0$. The induction hypothesis applied to M_1 , yields that $\text{Ext}_R^1(R/\mathfrak{a}, H_{\mathfrak{a}}^{t-1}(M_1))$ is \mathfrak{a} -minimax. Hence again by lemma 2.1, $\text{Ext}_R^1(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M))$ is \mathfrak{a} -minimax.

(ii) Let N be an \mathfrak{a} -minimax submodule of $H_{\mathfrak{a}}^t(M)$ and let L be a finitely generated R -module with $\text{Supp } L \subseteq V(\mathfrak{a})$. Hence by part (i) and [1, Theorem 4.2], the R -module $\text{Ext}_R^i(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax for all $i \leq 1$. Now, in view of [1, Theorem 2.7], the R -module $\text{Ext}_R^i(L, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax for all $i \leq 1$.

(iii) By the exact sequence

$$\text{Ext}_R^2(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M)) \rightarrow \text{Ext}_R^2(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M)/N) \rightarrow \text{Ext}_R^3(R/\mathfrak{a}, N)$$

and the fact that $\text{Ext}_R^3(R/\mathfrak{a}, N)$ is \mathfrak{a} -minimax, it is sufficient to show that the R -module $\text{Ext}_R^2(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M))$ is \mathfrak{a} -minimax. To do this, we use induction on t . When $t = 0$, by assumption $\text{Ext}_R^2(R/\mathfrak{a}, M)$ and $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^1(M))$ are both \mathfrak{a} -minimax. Therefore, by Proposition 2.2, the R -module $\text{Ext}_R^1(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M))$ is \mathfrak{a} -minimax. Thus the exact sequence

$$\text{Ext}_R^1(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M)) \rightarrow \text{Ext}_R^2(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(M)) \rightarrow \text{Ext}_R^2(R/\mathfrak{a}, M),$$

implies that $\text{Ext}_R^2(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(M))$ is \mathfrak{a} -minimax. Now suppose, inductively, that $t > 0$ and suppose that the result has been proved for $t - 1$. As $\Gamma_{\mathfrak{a}}(M)$ is \mathfrak{a} -cominimax, it follows from the exact sequence

$$\text{Ext}_R^{t+2}(R/\mathfrak{a}, M) \rightarrow \text{Ext}_R^{t+2}(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M)) \rightarrow \text{Ext}_R^{t+3}(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(M))$$

that $\text{Ext}_R^{t+2}(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M))$ is \mathfrak{a} -minimax. Therefore we may assume that M is \mathfrak{a} -torsion free. Let E be an injective envelope of M and put $M_1 = E/M$. By using a similar proof as in the proof of part (i), the induction hypothesis applied to M_1 , yields that $\text{Ext}_R^2(R/\mathfrak{a}, H_{\mathfrak{a}}^{t-1}(M_1))$ is \mathfrak{a} -minimax, and so $\text{Ext}_R^2(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M))$ is \mathfrak{a} -minimax.

(iv) Let N be an \mathfrak{a} -minimax submodule of $H_{\mathfrak{a}}^t(M)$ and let L be a finitely generated R -module with $\text{Supp } L \subseteq V(\mathfrak{a})$. Hence by parts (ii) and (iii), the R -module $\text{Ext}_R^i(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax for all $i \leq 2$. Now, in view of [1, Theorem 2.7], the R -module $\text{Ext}_R^i(L, H_{\mathfrak{a}}^t(M)/N)$ is \mathfrak{a} -minimax for all $i \leq 2$.

(v) In view of [1, Theorem 2.7] it is enough to show that $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(M)/N)$ is \mathfrak{a} -minimax. To this end, by the exact sequence

$$\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(M)) \rightarrow \text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(M)/N) \rightarrow \text{Ext}_R^1(R/\mathfrak{a}, N)$$

and that $\text{Ext}_R^1(R/\mathfrak{a}, N)$ is \mathfrak{a} -minimax, it is sufficient to show that $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(M))$ is \mathfrak{a} -minimax. To do this, we use induction on t . When $t = 0$, by assumption $\text{Ext}_R^1(R/\mathfrak{a}, M)$ and $\text{Ext}_R^2(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(M))$ are \mathfrak{a} -minimax, and so in view of Proposition 2.2 the R -module $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^1(M))$ is \mathfrak{a} -minimax. Now suppose, inductively, that $t > 0$ and suppose that the result has been proved for $t - 1$. Since $\Gamma_{\mathfrak{a}}(M)$ is \mathfrak{a} -cominimax, it follows from the exact sequence

$$\text{Ext}_R^{t+1}(R/\mathfrak{a}, M) \rightarrow \text{Ext}_R^{t+1}(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M)) \rightarrow \text{Ext}_R^{t+2}(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(M))$$

that $\text{Ext}_R^{t+1}(R/\mathfrak{a}, M/\Gamma_{\mathfrak{a}}(M))$ is \mathfrak{a} -minimax. Therefore we may assume that M is \mathfrak{a} -torsion free. Let E be an injective envelope of M and put $M_1 = E/M$. Again, by using a similar proof as in the proof of part (i), the induction hypothesis applied to M_1 , yields that $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^t(M_1))$ is \mathfrak{a} -minimax, and so $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(M))$ is \mathfrak{a} -minimax. \square

By using the above result, we can deduce the following corollary, which is the main result of this paper.

Corollary 2.4. *Let M be an \mathfrak{a} -minimax R -module, and let t be a non-negative integer such that $H_{\mathfrak{a}}^i(M)$ is \mathfrak{a} -cominimax for all $i < t$. Then the following hold:*

(i) *For any \mathfrak{a} -minimax submodule N of $H_{\mathfrak{a}}^t(M)$ and for any finitely generated R -module L with $\text{Supp } L \subseteq V(\mathfrak{a})$, the R -modules $\text{Hom}_R(L, H_{\mathfrak{a}}^t(M)/N)$ and $\text{Ext}_R^1(L, H_{\mathfrak{a}}^t(M)/N)$ are \mathfrak{a} -minimax.*

(ii) *If the R -modules $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(M))$ is \mathfrak{a} -minimax, then for any \mathfrak{a} -minimax submodule N of $H_{\mathfrak{a}}^t(M)$ and for any finitely generated R -module L with $\text{Supp } L \subseteq V(\mathfrak{a})$, the R -modules $\text{Hom}_R(L, H_{\mathfrak{a}}^t(M)/N)$, $\text{Ext}_R^1(L, H_{\mathfrak{a}}^t(M)/N)$ and $\text{Ext}_R^2(L, H_{\mathfrak{a}}^t(M)/N)$ are all \mathfrak{a} -minimax.*

Proof. Since M is \mathfrak{a} -minimax, so $\text{Ext}_R^i(R/\mathfrak{a}, M)$ is \mathfrak{a} -minimax for all i . Thus the result is followed by Theorem 2.3. \square

REFERENCES

- [1] J. Azami, R. Naghipour and B. Vakili, *Finiteness properties of local cohomology modules for \mathfrak{a} -minimax modules*, Proc. Amer. Math. Soc, **137** (2009), 439-448.
- [2] M.P. Brodmann and F.A. Lashgari, *A finiteness result for associated primes of local cohomology modules*, Proc. Amer. Math. Soc. **128** (2000). 2851-2853.
- [3] M. P. Brodmann and R. Y. Sharp, *Local cohomology: an algebraic introduction with geometric applications*, Cambridge University Press, 1998.
- [4] M. T. Dibaei and S. Yassemi, *Finiteness of extension functors of local cohomology modules*, Comm. in Algebra, **34** (2006), 3097-3101.
- [5] K. Divaani-Aazar and M.A. Esmkhani, *Artinianness of local cohomology modules of ZD -modules*, Comm. Algebra, **33** (2005), 2857-2863.
- [6] A. Grothendieck, *Local cohomology*, Lecture Notes in Mathematics, **41**, (Springer, Berlin, 1967). 161-166.
- [7] H. Zöschinger, *Minimax modulu*, J. Algebra, **102** (1986), 1-32.
- [8] H. Zöschinger, *Über die Maximalbedingung für radikalvolle Untermoduln*, Hokkaido Math. J. **17** (1988), 101-116.

DEPARTMENT OF MATHEMATICS, ISLAMIC AZAD UNIVERSITY, SHABESTAR BRANCH, IRAN.
E-mail address: bvakil2004@yahoo.com and bvakil@iaushab.ac.ir